

Identification of Diagnosis Fungi that Cause Potato Root Rot

Saja W. A'shour¹ and Theyab A. Farhan²

¹Department of Plant Protection, College of Agriculture, University of Anbar, Iraq.
saj20g7013@uoanbar.edu.iq.

²Department of Plant Protection, College of Agriculture, University of Anbar, Iraq.
deab.frahen@uoanbar.edu.iq

* Correspondence: saj20g7013@uoanbar.edu.iq.

Available from. <http://dx.doi.org/10.21931/BJ/2024.01.01.51>

ABSTRACT

Results of collecting samples from different regions of Anbar Governorate (Al-Amiriyah, Al-Khalidya, Fallujah, Heet and Ramadi) showed that potato root rot disease is widespread in all regions collected. The results of isolation and phenotypic and molecular diagnosis using the polymerase chain reaction (PCR) technique indicated showed the presence of fungus *Rhizoctonia* spp., and Fungus *Fusarium* spp. Accompanying potato root rot disease and the pathogenicity test using radish seeds on water Agar (W.R.) culture media, all tested isolates achieved a significant reduction in radish seed plants compared with control treatment uncontaminated by any of the isolates of fungi, which recorded infection rate 0%.

Keywords: Potato Root Rot, Diagnosis, Fungi, *Rhizoctonia solani*, *Fusarium solani*.

INTRODUCTION

Potato, *Solanum tuberosum* L., is considered a strategic vegetable crop and comes in fourth place after wheat, corn and rice. It is from the family *Solanaceae*, of strategic importance in most countries of the world in general and Iraq in particular, as it is considered one of the easiest crops to manage when compared to other vegetable crops as well as Most of its varieties, especially the early ones, complete their life cycle less than 100 days¹. Due to the ease of management of this crop and its high productivity and short life cycle, its cultivation has expanded significantly². As a result of the expansion in the cultivation of this crop, many problems appeared, foremost of which are fungal causes, especially root rot pathogens, leg ulcers and black crust, which cause economic losses. It increases farmers' production^{3, 4}. Root rot disease is one of the most essential plant diseases worldwide that affects many crops⁵. Root rot disease of potato plants caused by the fungi *Rhizoctonia solani* and *Fusarium solani* is among the most common diseases due to its highly destructive effects that may reach chronic effects in every potato growing area. Each of them can have an economic impact on the season and does not cause Economic losses are only on an annual basis and therefore considered a problem for producers who often suffer economic losses from this disease. Mold was first described as a potato tuber disease in Ireland in 1913, has been known to occur in North America since then, and has become a significant soil-borne disease. Statistics show that this species is becoming increasingly common, which poses a significant threat to potato production, especially in warm regions⁶. Root rot symptoms represent a

significant threat because the damage begins underground, where the first symptoms cannot be distinguished—the appearance of pathological symptoms on infected plants^{7, 8}. Because of the importance of the potato crop and the danger of root rot disease, the study aimed to isolate and characterize the phenotypic and molecular diagnosis of the fungi accompanying the samples infected with root rot disease.

MATERIALS AND METHODS

Sample collection

Samples were collected from potato plants that showed symptoms of disease (yellow, plant wilt, plant death, burn edges of leaves)⁹ from some fields of Anbar Governorate (Al-Khalidiya, Al-Amiriya, Sadat Al-Fallujah, Nuaimiya, Heet and Saqlawiyah) for the agricultural season Two thousand twenty-one for the spring loop, and the affected plant parts were placed in sterile polyethylene bags, with the place and date of sample collection recorded, and the samples were kept in the refrigerator at a temperature of 4 C° until the isolation process.

Isolation of fungi associated with potato root rot

Parts of roots that showed symptoms of rot were taken and washed with water to get rid of dust and cut into small pieces with a length of 0.5-1 cm. and superficially sterilized with sodium hypochlorite solution NaCl (chlorine 6% active substance) for 3 minutes after which the plant pieces were washed with distilled water Sterilized 3 times to get rid of the effects of the sterile material, dried with sterile filter paper and transferred by sterile forceps, and 4 plant pieces were planted in each Petri dish with a diameter of 9 cm containing the sterilized culture medium (PDA) Potato Dextrose Agar with an Autoclaved at 121 °C and a pressure of 5.1 bar/cm³ for 20 minutes and the antibiotic (Amoxicillin) was added to it at a concentration of 200 mg. L⁻¹, after sterilizing medium. The dishes were incubated at 25 ± 2 °C for 3 days, after which a microscopic examination was conducted to investigate the presence of pathogenic fungi.

Phenotypic diagnosis

Fungi isolates were diagnosed after the growth of fungal colonies on the PDA medium was prepared, as microscopic slides were examined using a light microscope, and depending on the cultural characteristics and using the taxonomic key specific to the species, the fungi were morphologically diagnosed¹⁰.

Molecular diagnosis

DNA was isolated and extracted in the Scientific Progress Laboratory / Baghdad - Al-Harithiya according to the ABIOPure protocol. Prepared by the Korean company Macrogen ITS13 -5'- TCCGTAGGTGAAC-CTGCGG, ITS43 - 5'-TCCTCCGCTTATTGATATGC. The Korean company Macrogen prepared the primers in lyophilized form in nuclease-free water to give a final concentration of 100 µl.µl⁻¹. A working solution of these primers was prepared by adding 10 µl of the starting primer (stored at -20 °C) to 90 µl of distilled water to obtain an effective starting solution of 10 µl. mol⁻¹ After PCR, gel electrophoresis to confirm the presence of amplification using polymerase chain reaction (PCR) was fully approved on the parameters of the extracted DNA. After completing the DNA amplification and migration steps on Agarose Gel, the samples were sent to the Korean Macrogen Company to obtain the sequence of nitrogenous bases (Sequencing). Several studies have relied on this technique in the diagnosis of fungi, as¹¹ showed that the PCR technique is used to diagnose *R. solani* fungi and relied in its diagnosis on the transcribed spacer internal region between (ITS4-ITS1) to distinguish between types of fungi. specialized.¹² also used PCR technique to diagnose species of the genus *Rhizoctonia* such as *R. solani*, isolated from different families of plants.

Pathogenicity tests of isolated fungi on radish seeds

The pathogenicity of seven isolates of *R.solani* spp. (Al-Amiriya - Rs1, Al-Amiriya 2 - Rs2, Al-Khalidiya - Rs3, Sadat Al-fallujah - Rs4, Al-Nuaimiya - Rs5, Heet - Rs6 and Al-Saqlawiyah- Rs7) And seven isolates of *Fusarium* spp. (Al-Amiriya - Fu1, Al-Amiriya 2 - Fu2, Al-Khalidiya - Fu3, Al-Khalidiya - Fu4, Al-Nuaimiya - Fu5, Heet - Fu6 Furthermore, Al-Saqlawiyah - Fu7) was tested. In the Department of Plant Protection at the College of Agriculture / University of Anbar, the test was carried out according to the method of ¹³ according to a Randomized completely design (CRD) sterile nutrient medium (Water Ager) was prepared and distributed in Petri dishes with a diameter of 9 cm containing 15-20 mm. From the more significant culture medium and water (W.A.) (20 g acre, liter of distilled water), sterilized by an autoclaved 121 °C and a pressure of 5.1 bar/cm³ for 20 minutes, added to the antibiotic Amoxicillin at a concentration of 200 mg/L⁻¹, the dishes were inoculated by taking a 5 mm diameter disc using A sterile cork piercing was placed in the middle of the dish. 3 dishes were used for each isolate of the tested fungi, and 3 were used as a comparison treatment (without adding a tablet from the mushroom colony). Then the dishes were transferred to the incubator and left at a temperature of 25 ± 2 for three days, after which the sterilized radish seeds were sown superficially with a solution of Sodium hypo chlori; 100%, 10 seeds were used for each plate. The seeds were sown in a circular motion around the fungal perennial disc so that the seeds were separated by 1 cm from the edge of the plate. Then, the plates were incubated for 14 days, after which the percentage of infection was calculated for each isolate according to the following equation:

$$\text{Germination \%} = \frac{\text{Number of Infected Plants}}{\text{Total Number of Plants}} \times 100.$$

RESULTS AND DISCUSSION

The results of sampling. in Fig. (1) and Fishow g. (1) Potato root rot disease is widespread in most potato growing areas in the world, and that is consistent with findings found by ^{14, 15}.



Figure 1: Potato plants infected with root rot disease.

No.	Sample Code	Location	Collection Date 2021
1	A1	Al-Amiriya	25 /10
2	A2	Al-Amiriya2	27 /10

3	A3	Al-Khalidiya	3 /11
4	A4	Sadat Al-Fallujah	5 /11
5	A5	Al-Nuaimiya	11 /11
6	A6	Heet	12 /11
7	A7	Al-Saqlawiyah	14 /12

Table 1: Areas from which samples were collected from Potato root rot disease.

Phenotypic diagnosis

The results of the phenotypic diagnosis shown in Table (2) showed the presence of the fungus *R.solani* spp. and *Fusarium* spp., which were associated with potato root rot disease, which is the main cause of the disease, and this is consistent with what was indicated by ^{16,17}.

No.	Sample Code	Fungi	Location	Collection Date 2021
1	A1	<i>Rhizoctoina</i> spp. and <i>Fusarium</i> spp.	Al-Amiriya	25 /10
2	A2	<i>Rhizoctoina</i> spp. and <i>Fusarium</i> spp.	Al-Amiriya 2	27 /10
3	A3	<i>Rhizoctoina</i> spp. and <i>Fusarium</i> spp.	Al-Khalidiya	3 /11
4	A4	<i>Rhizoctoina</i> spp. and <i>Fusarium</i> spp.	Sadat Al-Fallujah	5 /11
5	A5	<i>Rhizoctoina</i> spp. and <i>Fusarium</i> spp.	Al-Nuaimiya	11 /11
6	A6	<i>Rhizoctoina</i> spp. and <i>Fusarium</i> spp.	Heet	12 /11
7	A7	<i>Rhizoctoina</i> spp. and <i>Fusarium</i> spp.	Al-Saqlawiyah	14 /12

Table 2: The most important fungi related to potato root rot were isolated and diagnosed phenotypically to the genus level.

No.	Location	Code	Fungi	Accession
1	Al-Amiriya	A1	<i>Fusarium Falciforme</i>	MN545530.1
2	Al-Khalidiya	K	<i>Fusarium Oxysporum</i>	OL691079.1
3	Al-Saqlawiyah	B	<i>Rhizoctoina solani</i>	LC507904.1
4	Sadat Al-fallujah	C	<i>Rhizoctoina solani</i>	MG654436.1
5	Al-Nuaimiya	D	<i>Rhizoctoina solani</i>	KY965394.1
6	Heet	E	<i>Rhizoctoina solani</i>	KY965394.1
7	Al-Khalidiya	F	<i>Rhizoctoina solani</i>	AJ318420.1
8	Al-Amiriya 2	A2	<i>Fusarium Keratoplasticum</i>	KY965394.1
9	Heet	A3	<i>Rhizoctoina solani</i>	MF440630.1

Table 3: Fungi associated with potato root rot diagnosed by PCR technique.




Figure 2: Results of ITS gene amplification of fungal species on Agarose Gel.

The pathogenicity test of some fungi isolates and their effect on the germination of radish seeds on the culture medium Water agar (W.A.)

Results showed in Table (4) that all tested fungi isolates tested on radish seeds were isolates of *Rhizoctonia* spp. and isolates of *Fusarium* spp. It significantly reduced the germination of radish seeds on water agar (W.A.) culture media compared to the control treatment that was not contaminated with any of the tested isolates *Rhizoctonia*, in which the infection rate was 0%. The test also showed the variation of the isolates in their effect on the germination of radish seeds, and the most influential of them was the isolate *Rhizoctonia* taken from Al-Anbar, Heet region, in which the infection rate was 90%. In contrast, *Fusarium* isolate is the strongest in the Al-Saqlawiya region, which achieved a 90% infection rate. Results agreed with that mentioned, and y 20, 21.

Location	Isolation code <i>Rhizoctonia</i>	Infection ratio %	Isolation code <i>Fusarium</i>	Infection ratio %
Amiriyah	<i>Rhizoctonia</i> Rs1	83.334	<i>Fusarium</i> Fu1	83.334
Amiriyah 2	<i>Rhizoctonia</i> Rs2	80	<i>Fusarium</i> Fu2	86.667
Khalidiya	<i>Rhizoctonia</i> Rs3	80	<i>Fusarium</i> Fu3	80
Sadat Al-Fallujah	<i>Rhizoctonia</i> Rs4	76.667	<i>Fusarium</i> Fu4	83.334
Nuaimiya	<i>Rhizoctonia</i> Rs5	90	<i>Fusarium</i> Fu5	83.334
Heet	<i>Rhizoctonia</i> Rs6	76.667	<i>Fusarium</i> Fu6	86.667

Saqlawiyah	<i>Rhizoctonia</i> Rs7	83.334	<i>Fusarium</i> Fu7	90
	Control	0.00	Control	0.00
		LSD 0.817		LSD 1.236

Table 4: Effects of different isolates of *Rhizoctonia* spp. and *Fusarium* spp. in germination of Radish seeds on water agar medium.




Figure 3: Fungus *Rhizoctonia* spp. and *Fusarium* spp. in germination of radish seeds on Water Agar media.

Molecular Diagnostics

Table (3) shows the results of the molecular diagnosis that the fungi that cause potato root rot disease are of the genus *Rhizoctonia* spp. and *Fusarium* spp., according to the arrangement of nitrogenous bases in the single DNA strand. These results are consistent with what was mentioned by ^{16, 18}, and 19, and these fungi are the leading cause of potato root rot disease.

CONCLUSIONS

Potato root rot was observed as a disease in all sample collections. Two fungi, *Rhizoctonia* spp, mainly caused and *Fusarium* spp. according to the results of phenotypic and molecular diagnosis.

REFERENCES

1. Tunio M.H., Jianmin G., Sher A. S., Imran A. L., Waqar A. Q., Kashif A. S., and F. A. Chandio. 2020. Potato production in aeroponics: An emerging food growing system in sustainable agriculture for food security. Chilean Journal of Agricultural Research. 80: (1).
2. Nasir, M.W. and Toth, Z.2022. Effect of Drought Stress on Potato Production: A Review. Agronomy, 12, 635. <https://doi.org/10.3390/agronomy12030635>.

3. Kamar U. , Sami u., Shafi M. , Shakil A. P. , Razia S. , Amit H. , Chao W. and C. H. Jian. 2021. In vitro and silico approach of fungal growth inhibition by *Trichoderma asperellum* HbGT6-07 derived volatile organic compounds. *Arabian Journal of Chemistry*. Volume 14, Issue 9.
4. Vilvert E., Stridh L. , Andersson B., Olson K., Aldén L. and A. Berlin. 2022. Evidence-based disease control methods in potato production: a systematic map protocol. *Environmental Evidence* .11:6. <https://doi.org/10.1186/s13750-022-00259-x> .
5. Bodah E.T. 2017. Root Rot Diseases in Plants: A Review of Common Causal Agents and Management Strategies. *Agri Res & Tech: Open Access J* 5(3): ARTOAJ.MS.ID.555661.
6. Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J. 2016. FUN Guild is an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol.* 20, 241–248. doi: 10.1016/j.funeco. 2015.06.006.
7. Jiang, J., Yu, M., Hou, R., Li, L., Ren, X., Jiao, C. 2019. Changes in the soil microbial community are associated with *Panaxquin quefolius* L. root rot diseases. *Plant Soil* 438, 143–156. doi: 10.1007/s11104-018-03928-4.
8. Giachero, M. L.; Declerck, S.; Marquez, N. 2022. Phytophthora Root Rot: Importance of the Disease, Current and Novel Methods of Control. *Agronomy*, 12, 610. <https://doi.org/10.3390/agronomy12030610>.
9. Agrios , G. N. 2005. *Plant Pathology* . 5th Ed. Elsevier Inc. USA. 998 pp.
10. Barnett, H. L., and B. B.Hunter. 1998. *Illustrated genera of imperfect fungi*. The American Phytopathological Society. U.S. Department of Agriculture, Agricultural Research Service, Washington State University, Pullman. APS Press. USA. St. Paul, Minnesota USA. 218p.
11. Lee, Y.M., Y.K. Choi and B.R. Min. 2000. Molecular characterization of *Fusarium solani* and its form *aespeiales* based on sequences analysis of the internal transcribed spacer (ITS) region of ribosomal DNA. *Mycobiology*. 28(2): 82-88.
12. Alhussaini M.S , Moslem M.A , Alghonaim M.I., Al-Ghanayem A.A. , Al-Yahya A.A.I ,HefnyH.M. and A. M. Saadabi. 2016. Characterization of *Cladosporium* Species by Internal Transcribed Spacer - PCR and Microsatellites-PCR. *Pak. J. Biol Sci.*; 19(4): 143-157. doi: 10.3923/pjbs . 143.157.
13. Bolkan, H.H. and Butler, E.E. 1974. Studies on Hetero karyosis Virulence of *Rhizoctonia solani*. *Phytopathology*, 64: 513-522.
14. Vatankhah M. , Saberi -Riseh R. , Eskandari M. M. and H. A. fzali. 2019. Evaluation of some fungicides for the control of *Fusarium* dry rot of potato. *J. Crop Prot.* 2019, 8 (3): 275 -285 .
15. El-kazzaz, M.K.; Ghoneim, K.E.; Agha, M.K.M.; Helmy, A.; Behiry, S.I.; Abdelkhalek, A.; Saleem, M.H.; Al-Askar, A.A.; Arishi, A.A.; Elsharkawy, M.M. 2022. Suppression of Pepper Root Rot and Wilt Diseases Caused by *Rhizoctonia solani* and *Fusarium oxysporum*. *Life*, 12, 587. <https://doi.org/10.3390/life12040587> .
16. Errampalli, D. and Johnston, H. W. 2001. Control of tuber -born black scurf (*Rhizoctonia solani*) and common scab (*Streptomyces scabies*) of potato with a combination of sodium hypochlorite and thiophanate -methyl preplanting seed tuber treatment. *Canadian Journal of Plant Pathology*, 23: 68 -77.
17. Sandipan, P. B., Solanki, B. P., Patel, N. N., Patel, R. L., Verma, P. D. and Desai, H. R. 2016. Effect of different fungicides against dry rot pathogens of potato caused by *Fusarium* spp. under in vitro condition. *Cercetari Agronomic in Moldova*, 4 (168): 69 -74.

Publisher's Note: Bionatura Journal stays neutral concerning jurisdictional claims in published maps and institutional affiliations.

Copyright: © 2024 by the authors. They were submitted for possible open-access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).